Journal of Biochemical and Biophysical Methods, 16 (1988) 330–332
Elsevier

Author Index

Adamcová, J., see Baláž, Š. (16) 75
Adessi, G.L., see Jouvenot, M. (16) 301
Aki, H., see Yamamoto, M. (16) 271
Alemany, M., see Vendrell, M. (16) 49
Alemany, M., see Zamora, F. (16) 293
Andrasko, J., see Gåveby, B.-M. (16) 141
Ansorge, W. and Pepperkok, R.
Performance of an automated system for capillary microinjection into living cells (16) 283
Arnaud, P., see Miribel, L. (16) 1
Arola, L., see Zamora, F. (16) 293
Avila, J., see Hernández, M.A. (16) 17

Bacon, T.A., Morvan, F., Rayner, B., Imbach, J.-L. and Wickstrom, E.
α-Oligodeoxynucleotide stability in serum, subcellular extracts and culture media (16) 311
Baláž, Š., Kuchár, A., Dřevojánek, J., Adamcová, J. and Vrbanová, A.
Liposome/saline partition coefficients of low-molecular-weight solutes by gel chromatography (16) 75
Bankovskaya, S.A., see Rekharsky, M.V. (16) 319
Barnett, S.F., LeStourgeon, W.M. and Friedman, D.L.
Rapid purification of native C protein from nuclear ribonucleoprotein particles (16) 87
Barzaghi, B., see Rabilloud, T. (16) 237
Bassols, A.M., Carreras, J. and Cussó, R.
A method for determination of glucose 1,6-bisphosphatase (16) 55
Berggren, M., see Cotgreave, I.A. (16) 247
Bērziņa-Bērzīte, R.V., see Rekharsky, M.V. (16) 319
Bjellqvist, B., see Östergren, K. (16) 165
Bontempi, L., see Gelfi, C. (16) 193
Bossi, M.L., Bossi, O., Gelfi, C. and Righetti, P.G.
Casting immobilized pH gradients into cylindrical polyacrylamide gels (16) 171
Bossi, O., see Bossi, M.L. (16) 171
Buettner, G.R.
In the absence of catalytic metals ascorbate does not autoxidize at pH 7: ascorbate as a test for catalytic metals (16) 27

Cantisani, A.M., see Conti, A. (16) 205
Carreras, J., see Bassols, A.M. (16) 55
Cattò, N., see Gelfi, C. (16) 193
Celentano, F.C., Tonani, C., Fazio, M., Gianazza, E. and Righetti, P.G.
pH gradients generated by polyprotic buffers. I. Theory and computer simulation (16) 109
Celentano, F.C., see Righetti, P.G. (16) 129
Chiari, M., see Righetti, P.G. (16) 185
Conti, A., Napolitano, L., Cantisani, A.M., Davoli, R. and Dall'Olio, S.
Bovine β-lactoglobulin H: isolation by preparative isoelectric focusing in immobilized pH gradients and preliminary characterization (16) 205
Cotgreave, I.A., Weis, M., Berggren, M., Sandy, M.S. and Moldéus, P.W.
Determination of the intracellular protein thiol distribution of hepatocytes using monobromobimane derivatisation of intact cells and isolated subcellular fractions (16) 247
Cserháti, T., Szögyi, M. and Lelkes, L.
Reversed-phase thin-layer chromatography used to study the simultaneous effect of pH and ion strength on the lipophilicity of chlorhexidine (16) 263
Cussó, R., see Bassols, A.M. (16) 55

Dall'Olio, S., see Conti, A. (16) 205
Davoli, R., see Conti, A. (16) 205
Descotes, F., see Jouvenot, M. (16) 301
Dhainaut, F., see Pelletier, B. (16) 63
Domscheit, A., see Gåveby, B.-M. (16) 141
Dřevojánek, J., see Baláž, Š. (16) 75

Egorov, A.M., see Rekharsky, M.V. (16) 319
Eriksson, G., see Östergren, K. (16) 165

0165-022X/88/$03.50 © 1988 Elsevier Science Publishers B.V. (Biomedical Division)

Fazio, M., see Celentano, F.C. (16) 109
Fazio, M., see Righetti, P.G. (16) 129
Friedman, D.L., see Barnett, S.F. (16) 87
Fukal, L. and Kasafírek, E.
N-Succinyl-alanyl-methionyl-*S*-benzylcysteine *p*-nitroanilide as a sensitive substrate for assaying activity of cysteine proteinases (16) 323

Galchenko, G.L., see Rekharsky, M.V. (16) 319
Gåveby, B.-M., Pettersson, P., Andrasko, J., Ineva-Flygare, L., Johannesson, U., Görg, A., Postel, W., Domscheit, A., Mauri, P.L., Pietta, P., Gianazza, E. and Righetti, P.G.
Stable storage conditions of Immobiline chemicals for isoelectric focusing (16) 141
Gelfi, C., Righetti, P.G., Cattò, N., Bontempi, L. and Gianazza, E.
Immunoblotting from immobilized pH gradients. The case of α_2-macroglobulin (16) 193
Gelfi, C., see Bossi, M.L. (16) 171
Gianazza, E., see Celentano, F.C. (16) 109
Gianazza, E., see Gåveby, B.-M. (16) 141
Gianazza, E., see Gelfi, C. (16) 193
Gianazza, E., see Miribel, L. (16) 1
Gianazza, E., see Righetti, P.G. (16) 129
Görg, A., see Gåveby, B.-M. (16) 141
Guan, A.L. and Markland, Jr., F.S.
Isoelectric focusing in immobilized pH gradients of a snake venom fibrinolytic enzyme (16) 215

Hernández, M.A., Padilla, R. and Avila, J.
Iodination of proteins on nitrocellulose blotting paper (16) 17

Imbach, J.-L., see Bacon, T.A. (16) 311
Ineva-Flygare, L., see Gåveby, B.-M. (16) 141

Johannesson, U., see Gåveby, B.-M. (16) 141
Jouvenot, M., Descotes, F., Remy-Martin, A. and Adessi, G.L.
A sulfone group-labeled TEM-DNA probe: comparison with a ^{32}P-labeled probe in dot-hybridization (16) 301

Kasafírek, E., see Fukal, L. (16) 323
Khaleque, M.A. and Sawicki, C.A.
Photolysis method for determination of the tetramer-dimer dissociation constant of deoxyhemoglobin (16) 41
Kuchár, A., see Baláž, Š. (16) 75

Lelkes, L., see Cserháti, T. (16) 263
LeStourgeon, W.M., see Barnett, S.F. (16) 87
Lopatnev, S.V., see Rekharsky, M.V. (16) 319

Markland, Jr., F.S., see Guan, A.L. (16) 215
Mauri, P.L., see Gåveby, B.-M. (16) 141
Miribel, L., Gianazza, E. and Arnaud, P.
The use of dye-ligand affinity chromatography for the purification of non-enzymatic human plasma proteins (16) 1
Moldéus, P.W., see Cotgreave, I.A. (16) 247
Morvan, F., see Bacon, T.A. (16) 311

Napolitano, L., see Conti, A. (16) 205

Onyezili, F.N.
The enzyme coupling process in urease immobilization on *O*-alkylated nylon tubes (16) 255
Östergren, K., Eriksson, G. and Bjellqvist, B.
The influence of the support material used on band sharpness in Immobiline gels (16) 165
Ozoliņa, R.K., see Rekharsky, M.V. (16) 319

Padilla, R., see Hernández, M.A. (16) 17
Pauly, A., see Pelletier, B. (16) 63
Pelletier, B., Dhainaut, F., Pauly, A. and Zahnd, J.-P.
Evaluation of growth rate in adhering cell cultures using a simple colorimetric method (16) 63
Pepperkok, R., see Ansorge, W. (16) 283
Pernelle, J.-J., Righetti, P.G. and Wahrmann, J.P.
2-D analysis of human skeletal muscle myosin light chains with immobilized pH gradients (16) 227
Pettersson, P., see Gåveby, B.-M. (16) 141
Pietta, P., see Gåveby, B.-M. (16) 141
Postel, W., see Gåveby, B.-M. (16) 141

Rabilloud, T., Barzaghi, B. and Righetti, P.G.
Use of polybuffer as carrier ampholytes in mixed-bed Immobiline gels for isoelectric focusing (16) 237
Rayner, B., see Bacon, T.A. (16) 311
Rekharsky, M.V., Egorov, A.M., Bankovskaya, S.A., Ozoliņa, R.K., Vīna, I.A., Bērziņa-Bērzīte, R.V., Lopatnev, S.V. and Galchenko, G.L.
Microcalorimetric determination of substrate specificity of L-asparaginase in *Escherichia coli* and *Erwinia carotovora* (16) 319

Remy-Martin, A., see Jouvenot, M. (16) 301
Righetti, P.G.
Isoelectric focusing as the crow flies (16) 99
Righetti, P.G., Chiari, M., Sinha, P.K. and Santaniello, E.
Focusing of pepsin in strongly acidic immobilized pH gradients (16) 185
Righetti, P.G., Fazio, M., Tonani, C., Gianazza, E. and Celentano, F.C.
pH gradients generated by polyprotic buffers. II. Experimental validation (16) 129
Righetti, P.G., see Bossi, M.L. (16) 171
Righetti, P.G., see Celentano, F.C. (16) 109
Righetti, P.G., see Gåveby, B.-M. (16) 141
Righetti, P.G., see Gelfi, C. (16) 193
Righetti, P.G., see Pernelle, J.-J. (16) 227
Righetti, P.G., see Rabilloud, T. (16) 237

Sandy, M.S., see Cotgreave, I.A. (16) 247
Santaniello, E., see Righetti, P.G. (16) 185
Sawicki, C.A., see Khaleque, M.A. (16) 41
Sinha, P.K., see Righetti, P.G. (16) 185
Szögyi, M., see Cserháti, T. (16) 263

Tonani, C., see Celentano, F.C. (16) 109
Tonani, C., see Righetti, P.G. (16) 129

Vendrell, M. and Alemany, M.
An improved procedure for the isolation of rat brown adipose tissue cells (16) 49
Vīna, I.A., see Rekharsky, M.V. (16) 319
Vrbanová, A., see Baláž, Š. (16) 75

Wahrmann, J.P., see Pernelle, J.-J. (16) 227
Weis, M., see Cotgreave, I.A. (16) 247
Wickstrom, E., see Bacon, T.A. (16) 311

Yamamoto, M. and Aki, H.
Application of differential flow microcalorimetry for study of drug interactions in the blood system (16) 271

Zahnd, J.-P., see Pelletier, B. (16) 63
Zamora, F., Arola, L. and Alemany, M.
Some pitfalls and considerations of plasma ammonia estimation (16) 293

Journal of Biochemical and Biophysical Methods, 16 (1988) 333–334
Elsevier

Subject Index

Acidic pH range, (16), 185
Adrenergic stimulation, (16), 49
Alternative method, (16), 63
Anion-exchange chromatography, (16), 87
Ascorbate, (16), 27
Association constant by gel chromatography, (16), 75
Autoxidation, (16), 27
Band sharpness, (16), 165
Berthelot method, (16), 293
Binding heat, (16), 271
Brown adipocyte isolation, (16), 49
Brown adipose tissue, (16), 49
Buffering power, (16), 109; (16), 129
C protein, (16), 87
Carrier ampholyte, (16), 237
Cell culture, (16), 63
Centrifugation by rotation, (16), 171
Chlorhexidine, (16), 263
Chlorpromazine hydrochloride, (16), 271
Chromogenic substrate, (16), 323
Computer automatisation, (16), 283
Computer simulation, (16), 109
Copper, (16), 27
Cysteine proteinase activity, (16), 323
Deoxyhemoglobin dimerization, (16), 41
Deoxyribonuclease, (16), 311
Differential flow microcalorimeter, (16), 271
Dot-hybridization, (16), 301
Dye-ligand affinity chromatography, (16), 1
Enzyme coupling, (16), 255
Epithelial cell, (16), 63
Erythrocyte, (16), 271
Fibrinolytic isoenzyme, (16), 215
Fibroblast, (16), 63
FITC-dextran, (16), 283
Fluorescent label, (16), 283
Gel chromatography, (16), 75
Glass capillary, (16), 283
Glucose 1,6-bisphosphatase, (16), 55
Glutamate dehydrogenase method, (16), 293
Gram-negative bacteria, (16), 301
Growth rate evaluation, (16), 63
HeLa cell lysate, (16), 311
Hemolysis, (16), 271
High-performance liquid affinity chromatography, (16), 1
Immobiline gel, (16), 165
Immobilized pH gradient, (16), 99; (16), 109; (16), 129; (16), 141; (16), 171; (16), 185; (16), 193; (16), 205; (16), 215; (16), 227; (16), 237
Intact cell, (16), 247
Iodometric method, (16), 293
Ionic strength, (16), 109; (16), 129
Iron, (16), 27
Isoelectric focusing, (16), 99; (16), 141; (16), 165; (16), 171; (16), 185; (16), 193; (16), 205; (16), 215; (16), 227; (16), 237
β-Lactoglobulin H, (16), 205
Lipophilic character, (16), 263
Liposome, (16), 75
Liposome/saline partition coefficient, (16), 75
α_2-Macroglobulin, (16), 193
Metalloproteinase, (16), 215
Methylene blue, (16), 63
Microcalorimetry, (16), 319
Microinjection, (16), 283
Microtubule associated protein, (16), 17
Monobromobimane, (16), 247
Myosin, (16), 227
Nessler method, (16), 293
p-Nitroanilide, (16), 323
α-Oligodeoxynucleotide hydrolysis, (16), 311
Oxygen consumption, (16), 49
Papain, (16), 323
Pepsin, (16), 185
Peptide mapping, (16), 17
pH gradient, (16), 109
Photolysis, (16), 41
Plasma ammonia, (16), 293
Plasma protein, human, (16), 271
Plasma protein purification, human, (16), 1
Polyprotic buffer, (16), 109; (16), 129
Protein thiol analysis, (16), 247
Purification, (16), 87
Radioactive labeling, (16), 17

0165-022X/88/$03.50 © 1988 Elsevier Science Publishers B.V. (Biomedical Division)

Radioenzymatic method, (16), 293
Reticulocyte lysate, rabbit, (16), 311
Reversed-phase thin-layer chromatography, (16), 263
Ribonucleoprotein, (16), 87
Serum, fetal bovine, (16), 311
Serum, human, (16), 311
Snake venom, (16), 215
Solid-phase assay, (16), 17
Stable acrylamide derivative, (16), 141
Stepwise regression analysis, (16), 263
Subcellular fraction, (16), 247
Substrate specificity of enzymes, (16), 319
Sulfone-labeled DNA probe, (16), 301
Support material, (16), 165
TEM-type β-lactamase gene, (16), 301
Tetramer-dimer dissociation, (16), 41
Two-dimensional map, (16), 141; (16), 227
Urease immobilization, (16), 255